arXiv:1502.05510v3 [math.ST] 21 Jan 2016

Unbiased estimation of the volume of a convex body

Nikolay Baldin
Institute of Mathematics
Humboldt-Universität zu Berlin
baldin.np@gmail.com

Markus Reiß*
Institute of Mathematics
Humboldt-Universität zu Berlin
mreiss@math.hu-berlin.de

January 22, 2016

Abstract

Based on observations of points uniformly distributed over a convex set in $\mathbb{R}^d$, a new estimator for the volume of the convex set is proposed. The estimator is minimax optimal and also efficient non-asymptotically: it is nearly unbiased with minimal variance among all unbiased oracle-type estimators. Our approach is based on a Poisson point process model and as an ingredient, we prove that the convex hull is a sufficient and complete statistic. No hypotheses on the boundary of the convex set are imposed. In a numerical study, we show that the estimator outperforms earlier estimators for the volume. In addition, an adjusted set estimator for the convex body itself is proposed.

Keywords: volume estimation, convex hull, Poisson point process, UMVU, stopping set, exact oracle inequality, missing volume

MSC code: 60G55, 62G05, 62M30

*We are grateful for helpful comments and questions by Mathias Reitzner and the anynonymous referee. Financial support by the DFG via Research Unit FOR1735 *Structural Inference in Statistics* is gratefully acknowledged.

1 Introduction

Driven by applications in image analysis and signal processing, the estimation of the support of a density attracts a lot of statistical activity. In many cases it is natural to assume a convex shape for the support set. First fundamental results for convex support estimation have been achieved by Korostelev and Tsybakov (1993, 1994) who prove minimax-optimal rates of convergence in Hausdorff distance for a set estimator. In particular, Korostelev and Tsybakov (1993) prove that the convex hull of the points $\widehat{C}_n$, which is a maximum likelihood estimator for the set C, is rate-optimal. Interestingly, the volume $|\widehat{C}_n|$ of the convex hull is not rate-optimal for estimation of the volume $|C|$ of the convex set and an alternative two-step estimator, optimal up to a logarithmic factor, was proposed. A fully rate-optimal estimator for the volume of a convex set with smooth boundary was then constructed by Gayraud (1997) based on three-fold sample splitting. For various extensions and applications of convex support estimation, let us refer to Mammen and Tsybakov (1995); Guntuboyina (2012); Brunel (2014) and the literature cited there. Related ideas under Hölder and monotonicity constraints, respectively, have been adopted by Reiß and Selk (2015) for a one-sided regression model.

Our contribution is the construction of a very simple volume estimator which is not only rate-optimal over all convex sets without boundary restrictions, but even adaptive in the sense that it attains almost the parametric rate if the convex set is a polytope. Our approach is non-asymptotic and provides much more precise properties. The analysis is based on a Poisson point process (PPP) observation model with intensity $\lambda > 0$ on the convex set $C \subseteq \mathbb{R}^d$. We thus observe

$$X_1, ..., X_N \overset{i.i.d.}{\sim} U(C), \quad N \sim \text{Poiss}(\lambda|C|),$$

where $(X_n), N$ are independent, see Section 2 below for a concise introduction to the PPP model. Using Poissonisation and de-Poissonisation techniques, this model exhibits asymptotic properties like the uniform model, i.e. a sample of $n = \lambda|C|$ uniformly on C distributed random variables $X_1, \dots, X_n$. The beautiful geometry of the PPP model, however, allows for much more concise ideas and proofs, see also Meister and Reiß (2013) for connections between PPP and regression models with irregular error distributions. From an applied perspective, PPP models are often natural, e.g. for spatial count data of photons or other emissions.

For known intensity λ of the PPP, we construct in Section 3 an *oracle* estimator $\widehat{\vartheta}_{oracle}$. Theorem 3.2 shows that this estimator is UMVU (uniformly of minimum variance among unbiased estimators) and rate-optimal. To this end, moment bounds from stochastic geometry for the missing volume of the convex hull, obtained by Bárány and Larman (1988) and Dwyer (1988) are essential. Moreover, we derive results of independent interest: the convex hull $\widehat{C} = \mathrm{conv}\{X_1, \ldots, X_N\}$ forms a sufficient and complete statistic (Proposition 3.5) and the Poisson point process, conditionally on $\widehat{C}$, remains Poisson within its convex hull (Theorem 3.1).

For the more realistic case of unknown intensity λ, we analyse in Section 4 our final estimator

$$\widehat{\vartheta} \stackrel{\text{def}}{=} \frac{N+1}{N_\circ + 1} |\widehat{C}|\,,$$

where $N_\circ$ denotes the number of observed points in the interior of $\widehat{C}$. We are able to prove a sharp oracle inequality, comparing the risk of this estimator to that of $\widehat{\vartheta}_{oracle}$. Here, very recent and advanced results by Reitzner (2003); Pardon (2011); Beermann and Reitzner (2015) on the variance of the number of points N_∂ on the boundary of $\widehat{C}$ and the missing volume $|C \setminus \widehat{C}|$ are of key importance. This fascinating interplay between stochastic geometry and statistics prevails throughout the work.

The lower bound showing that $\widehat{\vartheta}$ is indeed minimax-optimal is proved in Theorem 3.4 by adopting the proof of the lower bound in the uniform model by Gayraud (1997). A small simulation study is presented in Section 5. Moreover, we propose to enlarge the convex hull set by the factor $((N+1)/(N_\circ+1))^{1/d}$ and we study its error as an estimator of the set C itself. The proof of Lemma 4.1 is deferred to the Appendix.

2 Digression on Poisson Point Processes

Most of the results and notation are adapted from Karr (1991). We fix a compact convex set $\mathbf{E}$ in $\mathbb{R}^d$ with non-empty interior as a state space and denote by $\mathcal{E}$ its Borel σ-algebra. We define the family of convex subsets $\mathbf{C} = \{C \subseteq \mathbf{E}, \text{convex}, \text{closed}\}$ (this implies that all sets in $\mathbf{C}$ are compact) and the family of compact subsets $\mathbf{K} = \{K \subseteq \mathbf{E}, \text{compact}\}$. It is natural to equip the space $\mathbf{C}$ (resp. $\mathbf{K}$) with the Hausdorff-metric d_H and its Borel σ-algebra $\mathfrak{B}_{\mathbf{C}}$ (resp. $\mathfrak{B}_{\mathbf{K}}$). Then $(\mathbf{C}, d_H)$ is a compact and thus separable

space and the mapping $(x_1, \ldots, x_k) \mapsto \text{conv}\{x_1, \ldots, x_k\}$, which generates the convex hull of points $x_i \in \mathbf{E}$, is continuous from $\mathbf{E}^k$ to $(\mathbf{C}, d_H)$.

On $(\mathbf{E}, \mathcal{E})$ we define the set of point measures $\mathbf{M} = \{m \text{ measure on } \mathcal{E} : m(A) \in \mathbb{N}, \ \forall A \in \mathcal{E}\}$ equipped with the σ-algebra $\mathcal{M} = \sigma(m \mapsto m(A), A \in \mathcal{E})$. Let $C_c^+(\mathbf{E})$ be the collection of continuous functions $\mathbf{E} \mapsto [0, \infty)$ with compact support. A useful topology for $\mathbf{M}$ is the *vague topology* which makes $\mathbf{M}$ a complete, separable metric space, cf. Section 3.4 in Resnick (2013). A sequence of point measures $m_n \in \mathbf{M}$ then converges vaguely to a limit $m \in \mathbf{M}$ if and only if $m_n[f] \to m[f]$ for all $f \in C_c^+(\mathbf{E})$ where $m[f] = \int_{\mathbf{E}} f dm$. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be an abstract probability space. We call a measurable mapping $\mathcal{N} : \Omega \to \mathbf{M}$ a Poisson point process (PPP) of intensity $\lambda > 0$ on $C \in \mathbf{C}$ if

- for any $A \in \mathcal{E}$, we have $\mathcal{N}(A) \sim \text{Poiss}\big(\lambda |A \cap C|\big)$, where $|A \cap C|$ denotes the Lebesgue measure of $A \cap C$;
- for all mutually disjoint sets $A_1, ..., A_n \in \mathcal{E}$, the random variables $\mathcal{N}(A_1), ..., \mathcal{N}(A_n)$ are independent.

For statistical inference, we assume the Poisson point process to be defined on a set of non zero Lebesgue measure, i.e. $|C| > 0$. A more constructive and intuitive representation of the PPP $\mathcal{N}$ is $\mathcal{N} = \sum_{i=1}^N \delta_{X_i}$ for $N \sim \text{Poiss}(\lambda |C|)$ and i.i.d. random variables (X_i), independent of N and distributed uniformly $\mathbb{P}(X_i \in A) = |A \cap C| / |C|$, so that $\mathcal{N}(A) = \sum_{i=1}^N \mathbf{1}(X_i \in A)$ for any $A \in \mathcal{E}$.

We consider the convex hull of the PPP points $\widehat{C} : \mathbf{M} \to \mathbf{C}$ defined by $\widehat{C}(\mathcal{N}) := \text{conv}\{X_1, ..., X_N\}$, which by the above continuity property of the convex hull is a random element with values in the Polish space $(\mathbf{C}, d_H)$, see also Davis, Mulrow, and Resnick (1987) for a detailed study of the continuity of the convex hull. For a short notation, we shall further write $\widehat{C}$ to denote the convex hull of the process $\mathcal{N}$. In the sequel, conditional expectations and probabilities with respect to $\widehat{C}$ are thus well defined. We can also evaluate the probability

$$\mathbb{P}_C\big(\widehat{C} \in A\big) = \sum_{k=0}^{\infty} \frac{\mathrm{e}^{-\lambda |C|} \lambda^k}{k!} \int_{C^k} \mathbf{1}(\text{conv}\{x_1, ..., x_k\} \in A) d(x_1, ..., x_k)$$

for $A \in \mathfrak{B}_{\mathbf{C}}$. Usually, we only write the subscript C or sometimes (C, λ) when different probability distributions are considered simultaneously. The

likelihood function $\frac{d\mathbb{P}_{C,\lambda}}{d\mathbb{P}_{\mathbf{E},\lambda_0}}$ for $C \in \mathbf{C}$ and $\lambda, \lambda_0 > 0$ is then given by

$$\frac{d\mathbb{P}_{C,\lambda}}{d\mathbb{P}_{\mathbf{E},\lambda_0}}(X_1, ..., X_N) = \mathrm{e}^{\lambda_0|\mathbf{E}|-\lambda|C|}(\lambda/\lambda_0)^N \mathbf{1}(\forall\, i = 1, ..., N :\ X_i \in C)$$
$$= \mathrm{e}^{\lambda_0|\mathbf{E}|-\lambda|C|}(\lambda/\lambda_0)^N \mathbf{1}(\widehat{C} \subseteq C)\,, \tag{2.1}$$

cf. Thm. 1.3 in Kutoyants (1998). For the last line, we have used that a point set is in C if and only if its convex hull is contained in C.

For the set-indexed process $(\mathcal{N}(K), K \in \mathbf{K})$ we define its natural set-indexed filtration

$$\mathcal{F}_K \stackrel{\text{def}}{=} \sigma(\{\mathcal{N}(U); U \subseteq K, U \in \mathbf{K}\})$$

for any $K \in \mathbf{K}$. The filtration $(\mathcal{F}_K, K \in \mathbf{K})$ possesses the following properties:

- monotonicity: $\mathcal{F}_{K_1} \subseteq \mathcal{F}_{K_2}$ for any $K_1, K_2 \in \mathbf{K}$ with $K_1 \subseteq K_2$,
- continuity from above: $\mathcal{F}_K = \cap_{i=1}^{\infty} \mathcal{F}_{K_i}$ if $K_i \downarrow K$;

cf. Zuyev (1999). By construction, the restriction $\mathcal{N}_K = \mathcal{N}(\cdot \cap K)$ of the point process $\mathcal{N}$ onto $K \in \mathbf{K}$ is $\mathcal{F}_K$-measurable (in fact, $\mathcal{F}_K = \sigma(\{\mathcal{N}_K(U); U \in \mathbf{K}\})$). In addition, it can be easily seen that $\mathcal{N}_K$ is a Poisson point process in $\mathbf{M}$, cf. the Restriction Theorem in Kingman (1992), and thus $\widehat{C}(\mathcal{N}_K) = \mathrm{conv}(\{X_1, \ldots, X_N\} \cap K)$ is by the above arguments $\mathcal{F}_K$-measurable.

A random compact set $\mathcal{K}$ is a measurable mapping $\mathcal{K} : (\mathbf{M}, \mathcal{M}) \to (\mathbf{K}, \mathfrak{B}_{\mathbf{K}})$. Note that Zuyev (1999) defines a random compact set as a measurable mapping from $(\mathbf{M}, \mathcal{M})$ to $(\mathbf{K}, \sigma_{\mathbf{K}})$ where $\sigma_{\mathbf{K}}$ is the so-called *Effros* σ-algebra generated by the sets $\{F \in \mathbf{K} : F \cap K \neq \emptyset\}$, $K \in \mathbf{K}$. Thanks to Thm. 2.7 in Molchanov (2006), the Effros σ-algebra $\sigma_{\mathbf{K}}$ induced on the family of compact sets $\mathbf{K}$ coincides with the Borel σ-algebra $\mathfrak{B}_{\mathbf{K}}$, and we prefer to stick to the first definition of a random compact set for convenience. Next, we recall the definition of stopping sets from Rozanov (1982) in complete analogy with stopping times.

Definition 2.1. *A random compact set $\mathcal{K}$ is called an $\mathcal{F}_K$-stopping set if $\{\mathcal{K} \subseteq K\} \in \mathcal{F}_K$ for all $K \in \mathbf{K}$. The sigma-algebra of $\mathcal{K}$-history is defined as $\mathcal{F}_{\mathcal{K}} = \{A \in \mathfrak{F} : A \cap \{\mathcal{K} \subseteq K\} \in \mathcal{F}_K \ \forall K \in \mathbf{K}\}$, where $\mathfrak{F} = \sigma(\mathcal{F}_K; K \in \mathbf{K})$.*

For a set $A \subseteq \mathbf{E}$ let A^c denote its complement.

Lemma 2.2. *The set $\widehat{\mathcal{K}} \stackrel{\text{def}}{=} \overline{\widehat{C}^c}$, the closure of the complement of the convex hull, is an $(\mathcal{F}_K)$-stopping set.*

Proof. We claim $\widehat{\mathcal{K}} \subseteq K$ if and only if $K^c \subseteq \operatorname{conv}(\{X_1, \dots, X_N\} \cap K)$. Indeed, if $\widehat{\mathcal{K}} \subseteq K$ holds, then the boundary $\partial\widehat{C} = \partial\widehat{\mathcal{K}}$ is in K which implies $\operatorname{conv}(\{X_1, \dots, X_N\} \cap K) = \widehat{C}$. Consequently, $K^c \subseteq \widehat{\mathcal{K}}^c \subseteq \widehat{C} = \operatorname{conv}(\{X_1, \dots, X_N\} \cap K)$ holds. Conversely, $K^c \subseteq \operatorname{conv}(\{X_1, \dots, X_N\} \cap K)$ implies immediately $K^c \subseteq \widehat{C}$ and thus $\widehat{C}^c \subseteq K$. Since K is closed, we obtain $\widehat{\mathcal{K}} \subseteq K$.

Since $\{X_1, \dots, X_N\} \cap K$ are the realisations of the point process inside K and the convex hull is measurable, we conclude $\{K^c \subseteq \operatorname{conv}(\{X_1, \dots, X_N\} \cap K)\} \in \mathcal{F}_K$. □

We shall further use the following short notation: $N = \mathcal{N}(C)$ denotes the total number of points, $N_\circ = \mathcal{N}(\widehat{C}^\circ)$ the number of points in the interior of the convex hull $\widehat{C}$ and $N_\partial = \mathcal{N}(\partial\widehat{C}) = \mathcal{N}(\partial\widehat{\mathcal{K}})$ the number of points on the boundary of the convex hull. For asymptotic bounds we write $f(x) = O(g(x))$ or $f(x) \lesssim g(x)$ if $f(x)$ is bounded by a constant multiple of $g(x)$ and $f(x) \sim g(x)$ if $f(x) \lesssim g(x)$ as well as $g(x) \lesssim f(x)$.

3 Oracle case: intensity λ is known

For a PPP on $C \in \mathbf{C}$ with intensity $\lambda > 0$, we know $N \sim \operatorname{Poiss}(\lambda|C|)$. In the oracle case, when the intensity λ is known, N/λ estimates $|C|$ without bias and yields the classical parametric rate in λ:

$$\mathbb{E}[(N/\lambda - |C|)^2] = \lambda^{-2} \operatorname{Var}(N) = \frac{|C|}{\lambda}.$$

Another natural idea might be to use the plug-in estimator $|\widehat{C}|$ whose error is given by the missing volume and satisfies

$$\mathbb{E}[(|\widehat{C}| - |C|)^2] = \mathbb{E}[|C \setminus \widehat{C}|^2] = O(|C|^{2(d-1)/(d+1)} \lambda^{-4/(d+1)}),$$

where the bound is obtained similarly to (3.4) and (3.5) below. This means that its error is of smaller order than λ^{-1} for $d \leqslant 2$, but larger for $d \geqslant 4$.

For any $d \geqslant 2$, however, both convergence rates are worse than the minimax-optimal rate $\lambda^{-(d+3)/(d+1)}$, established below.

The way to improve these estimators is to observe that by the likelihood representation (2.1) for $\lambda = \lambda_0$ and the Neyman factorisation criterion the convex hull is a sufficient statistic. Consequently, by the Rao-Blackwell theorem, the conditional expectation of N/λ given the convex hull $\widehat{C}$ is an estimator with smaller mean squared error (MSE).

The number of points N can be split into the number N_∂ of points on the boundary and the number $N_\circ$ of points in the interior of the convex hull. The following theorem is essential in deriving the oracle estimator. Although the statement of the theorem is quite intuitive and already used in Privault (2012), the proof turns out to be nontrivial and is deferred to the Appendix.

Theorem 3.1. *The number N_∂ of points on the boundary of the convex hull is measurable with respect to the sigma-algebra of $\widehat{\mathcal{K}}$-history $\mathcal{F}_{\widehat{\mathcal{K}}}$. The number of points in the interior of the convex hull $N_\circ$ is, conditionally on $\mathcal{F}_{\widehat{\mathcal{K}}}$, Poisson-distributed:*

$$N_\circ \,\big|\, \mathcal{F}_{\widehat{\mathcal{K}}} \sim \mathit{Poiss}(\lambda_\circ) \text{ with } \lambda_\circ \stackrel{\text{def}}{=} \lambda|\widehat{C}|. \tag{3.1}$$

In addition, we have $\mathcal{F}_{\widehat{\mathcal{K}}} = \sigma(\widehat{C})$, where the latter is the sigma-algebra $\sigma(\{\widehat{C} \subseteq B, B \in \mathbf{C}\})$ completed with the null sets in $\mathfrak{F}$.

With Theorem 3.1 at hand, we obtain the *oracle* estimator

$$\widehat{\vartheta}_{oracle} \stackrel{\text{def}}{=} \mathbb{E}\Big[\frac{N}{\lambda} \,\Big|\, \widehat{C}\Big] = \mathbb{E}\Big[\frac{N_\circ + N_\partial}{\lambda} \,\Big|\, \widehat{C}\Big] = |\widehat{C}| + \frac{N_\partial}{\lambda}\,, \tag{3.2}$$

where conditioning on $\widehat{C}$ means conditioning on $\sigma(\widehat{C}) = \mathcal{F}_{\widehat{\mathcal{K}}}$.

Theorem 3.2. *For known intensity $\lambda > 0$, the oracle estimator $\widehat{\vartheta}_{oracle}$ is unbiased and of minimal variance among all unbiased estimators (UMVU). It satisfies*

$$\mathrm{Var}(\widehat{\vartheta}_{oracle}) = \frac{1}{\lambda}\mathbb{E}[|C \setminus \widehat{C}|]\,.$$

Its worst case mean squared error over $\mathbf{C}$ decays as $\lambda \uparrow \infty$ like $\lambda^{-(d+3)/(d+1)}$ in dimension d:

$$\limsup_{\lambda\to\infty} \lambda^{(d+3)/(d+1)} \sup_{C\in\mathbf{C},|C|>0} \Big\{ |C|^{-(d-1)/(d+1)} \mathbb{E}\big[(\widehat{\vartheta}_{oracle} - |C|)^2\big] \Big\} < \infty\,.$$

Remark 3.3. *The theorem implies that the rate of convergence for the RMSE (root mean-squared error) of the estimator* $\widehat{\vartheta}_{oracle}$ *is* $\lambda^{-(d+3)/(2d+2)}$. *In Theorem 3.4 below, we prove that the lower bound on the minimax risk in the PPP model is of the same order implying that the rate is minimax-optimal. Even more, the oracle estimator is* adaptive *in the sense, that its rate is faster if the missing volume decays faster. In particular, for polytopes* C *it is shown in Bárány and Larman (1988) and independently in Dwyer (1988) that* $\mathbb{E}[|C \setminus \widehat{C}|] \sim \lambda^{-1}(\log(\lambda|C|))^{d-1}$, *which implies a faster (almost parametric) rate of convergence for the RMSE of the oracle estimator.*

Proof. The unbiasedness follows immediately from the definition (3.2). By the law of total variance, we obtain

$$\begin{aligned}\mathrm{Var}(\widehat{\vartheta}_{oracle}) &= \mathrm{Var}\left(\frac{N}{\lambda}\right) - \mathbb{E}\Big[\mathrm{Var}\left(\frac{N}{\lambda} \,|\, \widehat{C}\right)\Big] = \frac{|C|}{\lambda} - \mathbb{E}\Big[\mathrm{Var}\left(\frac{N_\circ}{\lambda} \,|\, \widehat{C}\right)\Big] \\ &= \frac{|C|}{\lambda} - \mathbb{E}\Big[\frac{\lambda_\circ}{\lambda^2}\Big] = \frac{1}{\lambda}\mathbb{E}[|C \setminus \widehat{C}|]\,. \end{aligned} \tag{3.3}$$

Proposition 3.5 below affirms that the convex hull $\widehat{C}$ is not only a sufficient, but also a complete statistic such that by the Lehmann-Scheffé theorem, the estimator $\widehat{\vartheta}_{oracle}$ has the UMVU property.

Finally, we bound the expectation of the missing volume $|C \setminus \widehat{C}|$ by Poissonisation, i.e. using that the convex hull $\widehat{C}$ in the PPP model conditionally on the event $\{N = k\}$ is distributed as the convex hull $\widehat{C}_k = \mathrm{conv}\{X_1, ..., X_k\}$ in the model with k uniform observations on C, for which the following upper bound is known (e.g., Bárány and Larman (1988)):

$$\sup_{C\in\mathbf{C}, |C|>0} \mathbb{E}\Big[\frac{|C \setminus \widehat{C}_k|}{|C|}\Big] = O\big(k^{-2/(d+1)}\big)\,. \tag{3.4}$$

Thus, it follows by a Poisson moment bound

$$\begin{aligned}\sup_{C\in\mathbf{C}, |C|>0} \mathbb{E}\Big[\frac{|C \setminus \widehat{C}|}{|C|^{(d-1)/(d+1)}}\Big] &= \sup_{C\in\mathbf{C}, |C|>0} \sum_{k=0}^{\infty} \frac{\mathrm{e}^{-\lambda|C|}(\lambda|C|)^k}{|C|^{-2/(d+1)}k!}\mathbb{E}\Big[\frac{|C \setminus \widehat{C}_k|}{|C|}\Big] \\ &= O\big(\lambda^{-2/(d+1)}\big)\,. \end{aligned} \tag{3.5}$$

This bound, together with (3.3), yields the assertion. □

The lower bound for the risk in the PPP framework can be derived from the lower bound in the uniform model with a fixed number of observations, see Thm. 6 in Gayraud (1997).

Theorem 3.4. *For estimating $|C|$ in the PPP model with parameter class $\mathbf{C}$, the following asymptotic lower bound holds*

$$\liminf_{\lambda\to\infty} \lambda^{(d+3)/(d+1)} \inf_{\widehat{\vartheta}_\lambda} \sup_{C\in\mathbf{C}} \mathbb{E}_C[(|C| - \widehat{\vartheta}_\lambda)^2] > 0\,, \tag{3.6}$$

where the infimum extends over all estimators $\widehat{\vartheta}_\lambda$ in the PPP model with intensity λ.

Proof. We use that an estimator $\widehat{\vartheta}_\lambda$ in the PPP model is an estimator in the uniform model on the event $\{N = n\}$. Then, due to the lower bound in the uniform model in Gayraud (1997), for a constant $c > 0$ and for all $n \in \mathbb{N}$ there exists a set $C_n \in \mathbf{C}$ with $|C_n| \sim 1$ such that for all $k = 1, ..., n$,

$$\mathbb{E}_{C_n}\big[(|C_n| - \widehat{\vartheta}_\lambda)^2 \,\big|\, N = k\big] > cn^{-(d+3)/(d+1)}, \quad a.s.$$

Then, in the PPP model for $C = C_{\lfloor\lambda\rfloor}$ with $\lambda|C| \geqslant 1$, we have

$$\begin{aligned}
\mathbb{E}_C[(|C| - \widehat{\vartheta}_\lambda)^2] &= \sum_{k\in\mathbb{N}} \mathbb{E}_C\big[(|C| - \widehat{\vartheta}_\lambda)^2 \,\big|\, N = k\big]\, \mathbb{P}(N = k) \\
&\geqslant \sum_{k\leqslant\lfloor\lambda\rfloor} \mathbb{E}_C\big[(|C| - \widehat{\vartheta}_\lambda)^2 \,\big|\, N = k\big]\, \mathbb{P}(N = k) \\
&> c\lfloor\lambda\rfloor^{-(d+3)/(d+1)} \big(1 - \mathbb{P}(N > \lfloor\lambda\rfloor)\big) \\
&\sim \lambda^{-(d+3)/(d+1)}\,,
\end{aligned}$$

applying Chernoff's inequality to $N \sim \mathrm{Poiss}(\lambda|C|)$ for the last line. Thus, the lower bound (3.6) follows. □

Proposition 3.5. *For known intensity $\lambda > 0$, the convex hull $\widehat{C} = conv\{X_1, ..., X_N\}$ is a complete statistic.*

Proof. We need to show the implication

$$\forall C \in \mathbf{C} : \mathbb{E}_C\big[T(\widehat{C})\big] = 0 \implies T(\widehat{C}) = 0 \quad \mathbb{P}_{\mathbf{E}} - a.s.$$

for any $\mathfrak{B}_{\mathbf{C}}$-measurable function $T : \mathbf{C} \to \mathbb{R}$. From the likelihood in (2.1) for $\lambda = \lambda_0$, we derive

$$\mathbb{E}_C\big[T(\widehat{C})\big] = \mathbb{E}_{\mathbf{E}}\Big[T(\widehat{C})\exp\big(\lambda|\mathbf{E}\setminus C|\big)\mathbf{1}(\widehat{C}\subseteq C)\Big]\,.$$

Since $\exp(\lambda|\mathbf{E}\setminus C|)$ is deterministic, $\mathbb{E}_C\big[T(\widehat{C})\big] = 0$ for all $C \in \mathbf{C}$ implies

$$\forall C \in \mathbf{C} : \mathbb{E}_{\mathbf{E}}\big[T(\widehat{C})\mathbf{1}(\widehat{C}\subseteq C)\big] = 0\,.$$

For $C \in \mathbf{C}$, define the family of convex subsets of C as $[C] = \{A \in \mathbf{C} | A \subseteq C\}$ such that $\widehat{C} \subseteq C \iff \widehat{C} \in [C]$. Splitting $T = T^+ - T^-$ with non-negative $\mathfrak{B}_{\mathbf{C}}$-measurable functions T^+ and T^-, we infer that the measures $\mu^{\pm}(B) = \mathbb{E}_{\mathbf{E}}[T^{\pm}(\widehat{C})\mathbf{1}(\widehat{C}\in B)]$, $B \in \mathfrak{B}_{\mathbf{C}}$, agree on $\{[C]\,|\,C \in \mathbf{C}\}$.

Note that the brackets $\{[C]|C \in \mathbf{C}\}$ are $\cap$-stable due to $[A]\cap[C] = [A\cap C]$ and $A\cap C \in \mathbf{C}$. If the σ-algebra $\mathcal{C}$ generated by $\{[C]\,|\,C\in\mathbf{C}\}$ contains $\mathfrak{B}_{\mathbf{C}}$, the uniqueness theorem asserts that the measures μ^+, μ^- agree on all Borel sets in $\mathfrak{B}_{\mathbf{C}}$, in particular on $\{T > 0\}$ and $\{T < 0\}$, which entails $\mathbb{E}_{\mathbf{E}}[T^+(\widehat{C})] = \mathbb{E}_{\mathbf{E}}[T^-(\widehat{C})] = 0$. Thus, in this case, $T(\widehat{C}) = 0$ holds $\mathbb{P}_{\mathbf{E}}$-a.s.

It remains to show that $\mathcal{C} = \sigma([C],\, C\in\mathbf{C})$ equals the Borel σ-algebra $\mathfrak{B}_{\mathbf{C}}$. This can be derived as a non-trivial consequence of Choquet's theorem, see Thm. 7.8 in Molchanov (2006), but we propose a short self-contained proof here. Let us define the family $\langle C\rangle = \{B \in \mathbf{C} | C \subseteq B\}$ of convex sets containing C. Then the closed Hausdorff ball with center C and radius $\varepsilon > 0$ has the representation

$$B_\varepsilon(C) \stackrel{\text{def}}{=} \{A\in\mathbf{C}\,|\,d_H(A,C)\leqslant\varepsilon\} = \{A\in\mathbf{C}\,|\,U_{-\varepsilon}(C)\subseteq A\subseteq U_\varepsilon(C)\}\,,$$

with $U_\varepsilon(C) = \{x\in\mathbf{E}\,|\,\mathrm{dist}(x,C)\leqslant\varepsilon\}$, $U_{-\varepsilon}(C) = \{x\in C\,|\,\mathrm{dist}(x,\mathbf{E}\setminus C)\leqslant\varepsilon\}$. Noting that $U_\varepsilon(C), U_{-\varepsilon}(C)$ are closed and convex and thus in $\mathbf{C}$, we obtain

$$B_\varepsilon(C) = \langle U_{-\varepsilon}(C)\rangle \cap [U_\varepsilon(C)]\,.$$

Since $(\mathbf{C}, d_H)$ is separable, our problem is reduced to proving that all angle sets $\langle C\rangle$ for $C\in\mathbf{C}$ are in $\mathcal{C}$. A further reduction is achieved by noting $\langle C\rangle = \bigcap_{x\in C}\langle x\rangle = \bigcap_{x\in C\cap\mathbb{Q}^d}\langle x\rangle$ setting $\langle x\rangle = \langle\{x\}\rangle$ for short such that it suffices to prove $\langle x\rangle\in\mathcal{C}$ for all $x\in\mathbf{E}$.

Now, let $x\in\mathbf{E}$ and $C\in\mathbf{C}$ such that $x\notin C$. Then, by the Hahn-Banach theorem, there are $\delta > 0, v\in\mathbb{R}^d$ such that $\langle v, c-x\rangle \geqslant \delta$ holds



Figure 1: The construction used in the proof.

for all $c \in C$. By a density argument, we may choose $\delta \in \mathbb{Q}^+$ and $v \in \mathbb{Q}^d$. Denoting the corresponding hyperplane intersected with $\mathbf{E}$ by $H_{\delta,v} = \{\xi \in \mathbf{E} \,|\, \langle v, \xi - x\rangle \geqslant \delta\}$, see Figure 1, we conclude

$$\langle x\rangle^{\complement} = \bigcup_{\delta\in\mathbb{Q}_+} \bigcup_{v\in\mathbb{Q}^d} \underbrace{[H_{\delta,v}]}_{\in\mathcal{C}} \in \mathcal{C}\,.$$

Consequently, $\langle x\rangle \in \mathcal{C}$ and thus $\mathfrak{B}_{\mathbf{C}} \subseteq \mathcal{C}$ hold. □

4 Unknown intensity λ: nearly unbiased estimation

In case the intensity λ is unknown and the oracle estimator $\widehat{\vartheta}_{oracle}$ in (3.2) is inaccessible, the maximum-likelihood approach suggests to use $N/|\widehat{C}|$ as an estimator for λ in (2.1). This yields the *plug-in* estimator for the volume,

$$\widehat{\vartheta}_{plugin} \stackrel{\text{def}}{=} |\widehat{C}| + \frac{N_\partial}{N}|\widehat{C}|\,.$$

In the unlikely event $N = |\widehat{C}| = 0$, we define $\widehat{\vartheta}_{plugin} = 0$. This estimator has a significant bias due to the following result, which is proved in the appendix.

Lemma 4.1. *For the bias of the plug-in MLE estimator $\widehat{\vartheta}_{plugin}$, it follows with some universal constant $c > 0$*

$$|C| - \mathbb{E}[\widehat{\vartheta}_{plugin}] \geqslant c\mathbb{E}[|\widehat{C} \setminus C|]^2\,, \quad \forall C \in \mathbf{C}\,. \tag{4.1}$$

The maximal bias over $C \in \mathbf{C}$ is thus at least of order $\lambda^{-4/(d+1)}$, which is worse than the minimax rate $\lambda^{-(d+3)/(2d+2)}$ for $d > 5$. Yet, in the two-dimensional finite sample study of Section 5 below, its performance is quite convincing. We surmise that $\widehat{\vartheta}_{plugin}$ is rate-optimal for $d \leqslant 5$, but we leave that question aside because the final estimator we propose will be nearly unbiased and will satisfy an *exact* oracle inequality. In particular, it is rate-optimal in any dimension. The new idea is to exploit that the number of interior points of $\widehat{C}$ satisfies $N_\circ \,\big|\, \widehat{C} \sim \mathrm{Poiss}(\lambda_\circ)$, see (3.1).

Remark 4.2. *There is no conditionally unbiased estimator for $\lambda_\circ^{-1}$ based on observing $N_\circ \,\big|\, \widehat{C} \sim Poiss(\lambda_\circ)$ for $\lambda_\circ$ ranging over some open (non-empty) interval. Otherwise, an estimator $\widetilde{\mu}(N_\circ)$ for $\lambda_\circ^{-1}$ would satisfy $\mathbb{E}[\widetilde{\mu}(N_\circ)|\widehat{C}] = \lambda_\circ^{-1}$ implying*

$$\sum_{k=0}^{\infty} \frac{\lambda_\circ^k}{k!} \widetilde{\mu}(k) \mathrm{e}^{-\lambda_\circ} = \lambda_\circ^{-1} \Rightarrow \sum_{k=0}^{\infty} \frac{\lambda_\circ^{k+1}}{k!} \widetilde{\mu}(k) = \sum_{k=0}^{\infty} \frac{\lambda_\circ^k}{k!} \,.$$

The coefficient for the constant term in the left and right power series would thus differ (0 versus 1), in contradiction with the uniqueness theorem for power series.

We provide an almost unbiased estimator for $\lambda_\circ^{-1}$ by noting that the first jump time of a time-indexed Poisson process with intensity ν is $\mathrm{Exp}(\nu)$-distributed and thus has expectation ν^{-1}. Taking conditional expectation of the first jump time with respect to the value of the Poisson process at time 1, we conclude that

$$\widehat{\mu}(N_\circ, \lambda_\circ) \stackrel{\mathrm{def}}{=} \begin{cases} (N_\circ + 1)^{-1}, & \text{for } N_\circ \geqslant 1, \\ 1 + \lambda_\circ^{-1}, & \text{for } N_\circ = 0 \end{cases}$$

satisfies $\mathbb{E}[\widehat{\mu}(N_\circ, \lambda_\circ)|\widehat{C}] = \lambda_\circ^{-1}$. Omitting the term $\lambda_\circ^{-1}$, depending on $\lambda_\circ$, in the unlikely case $N_\circ = 0$, we define our final estimator

$$\widehat{\vartheta} \stackrel{\mathrm{def}}{=} |\widehat{C}| + \frac{N_\partial}{N_\circ + 1} |\widehat{C}| \,.$$

For the proofs, we also define the *pseudo*-estimator

$$\widehat{\vartheta}_{pseudo} \stackrel{\mathrm{def}}{=} |\widehat{C}| + |\widehat{C}| N_\partial \Big(\frac{1}{N_\circ + 1} + \frac{\mathrm{e}^{-\lambda_\circ}}{\lambda_\circ} \Big) \,.$$

Theorem 4.3. *The pseudo-estimator* $\widehat{\vartheta}_{pseudo}$ *is unbiased and the estimator* $\widehat{\vartheta}$ *is asymptotically unbiased in the sense that with constants* $c_1, c_2 > 0$ *depending on* d, $d > 1$, *whenever* $\lambda|C| \geqslant 1$*:*

$$0 \leqslant |C| - \mathbb{E}[\widehat{\vartheta}] \leqslant c_1|C| \exp\bigl(-c_2(\lambda|C|)^{(d-1)/(d+1)}\bigr)\,, \quad \forall C \in \mathbf{C}\,.$$

Proof. We have

$$\mathbb{E}\Bigl[\frac{1}{N_\circ + 1} + \frac{\mathrm{e}^{-\lambda_\circ}}{\lambda_\circ} \,\Big|\, \widehat{C}\Bigr] = \mathrm{e}^{-\lambda_\circ}\lambda_\circ^{-1}\Bigl(\sum_{k=0}^{\infty} \frac{\lambda_\circ^{k+1}}{(k+1)k!} + 1\Bigr) = \lambda_\circ^{-1}\,,$$

which by $|\widehat{C}|\lambda_\circ^{-1} = \lambda^{-1}$ and $\mathbb{E}[\widehat{\vartheta}_{oracle}] = |C|$ implies unbiasedness of $\widehat{\vartheta}_{pseudo}$. Thus, it follows that

$$|C| - \mathbb{E}[\widehat{\vartheta}] = \mathbb{E}\bigl[|\widehat{C}|N_\partial \mathrm{e}^{-\lambda_\circ}\lambda_\circ^{-1}\bigr] = \lambda^{-1}\mathbb{E}\bigl[N_\partial \mathrm{e}^{-\lambda|\widehat{C}|}\bigr]\,.$$

We exploit the deviation inequality from Thm. 1 in Brunel (2013) and derive the bound for the exponential moment of the missing volume in the model with fixed number of points

$$\mathbb{E}[\exp\left(\lambda|C \setminus \widehat{C}_k|\right)] \leqslant b_1 \exp\left(b_2\lambda|C|k^{-2/(d+1)}\right), \quad k \geqslant 2\,,$$

for positive constants b_1, b_2, depending on the dimension according to Brunel (2013). For the cases $k = 0, 1$, we have the identity $\mathbb{E}[\exp\left(\lambda|C \setminus \widehat{C}_k|\right)] = \exp\left(\lambda|C|\right)$. By Poissonisation, similarly to (3.5), we derive

$$\exp(-\lambda|C|)\mathbb{E}[\exp\left(\lambda|C \setminus \widehat{C}|\right)] \leqslant b_3 \exp\bigl(-c_2(\lambda|C|)^{(d-1)/(d+1)}\bigr)\,, \quad (4.2)$$

for positive constants b_3, c_2, depending on the dimension. Hence, using the Cauchy-Schwarz inequality and the bound for the moments of the points on the convex hull,

$$\mathbb{E}[N_\partial^q] = O\bigl((\lambda|C|)^{q(d-1)/(d+1)}\bigr)\,, \quad q \in \mathbb{N}\,, \quad (4.3)$$

see e.g. Section 2.3.2 in Brunel (2014), we derive for a constant $c_1 > 0$

$$\begin{aligned}
\lambda^{-1}\mathbb{E}\bigl[N_\partial \mathrm{e}^{-\lambda|\widehat{C}|}\bigr] &\leqslant \lambda^{-1}\mathrm{e}^{-\lambda|C|}\mathbb{E}[N_\partial^2]^{1/2}\mathbb{E}[\mathrm{e}^{2\lambda|C\setminus\widehat{C}|}]^{1/2} \\
&\leqslant c_1\lambda^{-2/(d+1)}|C|^{(d-1)/(d+1)} \exp\bigl(-c_2(\lambda|C|)^{(d-1)/(d+1)}\bigr) \\
&\leqslant c_1|C| \exp\bigl(-c_2(\lambda|C|)^{(d-1)/(d+1)}\bigr)\,. \quad (4.4)
\end{aligned}$$

□

The next step of the analysis is to compare the variance of the pseudo-estimator $\widehat{\vartheta}_{pseudo}$ with the variance of the oracle estimator $\widehat{\vartheta}_{oracle}$, which is UMVU.

Theorem 4.4. *The following oracle inequality holds with a universal constant $c > 0$ and dimension-dependent constants $c_1, c_2 > 0$ for all $C \in \mathbf{C}$ with $\lambda|C| \geqslant 1$:*

$$\mathrm{Var}(\widehat{\vartheta}_{pseudo}) \leqslant (1 + c\alpha(\lambda, C))\,\mathrm{Var}(\widehat{\vartheta}_{oracle}) + r(\lambda, C)\,,$$

where

$$\begin{aligned}\alpha(\lambda, C) &= \frac{1}{|C|}\Big(\frac{1}{\lambda} + \frac{\mathrm{Var}(|C \setminus \widehat{C}|)}{\mathbb{E}[|C \setminus \widehat{C}|]} + \mathbb{E}[|C \setminus \widehat{C}|]\Big)\,,\\ r(\lambda, C) &= c_1(\lambda|C|)^{2(d-1)/(d+1)} \exp\big(-c_2(\lambda|C|)^{(d-1)/(d+1)}\big)\,.\end{aligned}$$

Proof. By the law of total variance, we obtain

$$\begin{aligned}\mathrm{Var}(\widehat{\vartheta}_{pseudo}) &= \mathrm{Var}\big(\mathbb{E}[\widehat{\vartheta}_{pseudo}|\widehat{C}]\big) + \mathbb{E}\big[\mathrm{Var}(\widehat{\vartheta}_{pseudo}|\widehat{C})\big]\\ &= \mathrm{Var}(\widehat{\vartheta}_{oracle}) + \mathbb{E}\Big[(N_\partial|\widehat{C}|)^2\,\mathrm{Var}\Big(\frac{1}{N_\circ + 1}|\widehat{C}\Big)\Big]\,.\end{aligned}$$

In view of $N_\circ \,|\, \widehat{C} \sim \mathrm{Poiss}(\lambda_\circ)$, a power series expansion gives

$$\mathbb{E}[(N_\circ + 1)^{-2}|\,\widehat{C}] = \lambda_\circ^{-1}\mathrm{e}^{-\lambda_\circ}\int_0^{\lambda_\circ}(\mathrm{e}^t - 1)/t\,dt\,.$$

The conditional variance can for $\lambda_\circ \to \infty$ thus be bounded by

$$\begin{aligned}\mathrm{Var}((1 + N_\circ)^{-1}|\,\widehat{C}) &\leqslant \lambda_\circ^{-1}\mathrm{e}^{-\lambda_\circ}\int_{\lambda_\circ/2}^{\lambda_\circ}\mathrm{e}^t/t\,dt - (\lambda_\circ)^{-2} + O(\mathrm{e}^{-\lambda_\circ/4})\\ &= (\lambda_\circ)^{-1}\int_0^{\lambda_\circ/2}\mathrm{e}^{-s}\Big(\frac{1}{\lambda_\circ - s} - \frac{1}{\lambda_\circ}\Big)\,ds + O(\mathrm{e}^{-\lambda_\circ/4})\\ &= \lambda_\circ^{-3}(1 + o(1))\,,\end{aligned}$$

where we have used $(\lambda_\circ - s)^{-1} - \lambda_\circ^{-1} = s\lambda_\circ^{-1}(\lambda_\circ - s)^{-1}$, $\int_0^\infty s\mathrm{e}^{-s}ds = 1$ and dominated convergence. Thanks to $(N_\circ + 1)^{-1} \in [0, 1]$ we conclude for some constant $c \geqslant 1$

$$\mathrm{Var}((1 + N_\circ)^{-1}|\,\widehat{C}) \leqslant c(1 \wedge \lambda_\circ^{-3}).$$

Consequently, we have

$$\begin{aligned}\mathrm{Var}(\widehat{\vartheta}_{pseudo}) &\leqslant \mathrm{Var}(\widehat{\vartheta}_{oracle}) + \mathbb{E}\big[(N_\partial|\widehat{C}|)^2 c(1 \wedge (\lambda|\widehat{C}|)^{-3})\big] \\ &= \mathrm{Var}(\widehat{\vartheta}_{oracle}) + c\mathbb{E}\big[(N_\partial|\widehat{C}|)^2 \wedge \lambda^{-3}(N_\partial)^2|\widehat{C}|^{-1}\big]\,,\end{aligned}$$

and with (3.3)

$$\begin{aligned}\frac{\mathrm{Var}(\widehat{\vartheta}_{pseudo})}{\mathrm{Var}(\widehat{\vartheta}_{oracle})} &\leqslant 1 + c\frac{\mathbb{E}\big[(N_\partial\lambda|\widehat{C}|)^2 \wedge (N_\partial)^2(\lambda|\widehat{C}|)^{-1}\big]}{\lambda\mathbb{E}[|C \setminus \widehat{C}|]} \\ &= 1 + c\frac{\mathbb{E}\big[(N_\partial)^2\big((\lambda|\widehat{C}|)^2 \wedge (\lambda|\widehat{C}|)^{-1}\big)\big]}{\mathbb{E}[N_\partial]}\,. \qquad (4.5)\end{aligned}$$

Define the 'good' event $\mathcal{G} = \{|\widehat{C}| > |C|/2\}$, on which $\big((\lambda|\widehat{C}|)^2 \wedge (\lambda|\widehat{C}|)^{-1}\big) \leqslant 2(\lambda|C|)^{-1}$. On the complement $\mathcal{G}^c$, we infer from $A^2 \wedge A^{-1} \leqslant 1$ for $A > 0$

$$\begin{aligned}\mathbb{E}\Big[(N_\partial)^2\big((\lambda|\widehat{C}|)^2 \wedge (\lambda|\widehat{C}|)^{-1}\big)\mathbf{1}_{\mathcal{G}^c}\Big] &\leqslant \mathbb{E}\big[N_\partial^2\mathbf{1}_{\mathcal{G}^c}\big] \\ &\leqslant \mathbb{E}[N_\partial^4]^{1/2}\mathbb{P}(|C \setminus \widehat{C}| \geqslant |C|/2)^{1/2} \\ &\leqslant c_1(\lambda|C|)^{2(d-1)/(d+1)} \exp\big(-c_2(\lambda|C|)^{(d-1)/(d+1)}\big)\,, \qquad (4.6)\end{aligned}$$

for some positive constant c_1 and c_2, using (4.2) and (4.3). It remains to estimate the upper bound (4.5) on $\mathcal{G}$

$$\frac{2c}{\lambda|C|}\frac{\mathbb{E}[N_\partial^2]}{\mathbb{E}[N_\partial]} = \frac{2c}{\lambda|C|}\Big(\frac{\mathrm{Var}(N_\partial)}{\mathbb{E}[N_\partial]} + \mathbb{E}[N_\partial]\Big)\,. \qquad (4.7)$$

Using the identity (17) in Beermann and Reitzner (2015) for the factorial moments for the number of vertices N_∂, we derive $\mathrm{Var}(N_\partial) \leqslant \lambda^2\,\mathrm{Var}(|C \setminus \widehat{C}|) + \lambda\mathbb{E}[|C \setminus \widehat{C}|]$ in view of $\mathbb{E}[N_\partial] = \lambda\mathbb{E}[|C \setminus \widehat{C}|]$. Thus, (4.7) is bounded by

$$\frac{2c}{\lambda|C|}\frac{\mathbb{E}[N_\partial^2]}{\mathbb{E}[N_\partial]} \leqslant \frac{2c}{|C|}\Big(\frac{1}{\lambda} + \frac{\mathrm{Var}(|C \setminus \widehat{C}|)}{\mathbb{E}[|C \setminus \widehat{C}|]} + \mathbb{E}[|C \setminus \widehat{C}|]\Big)\,,$$

which yields the assertion. □

As a result, we obtain an *oracle inequality* for the estimator $\widehat{\vartheta}$.

Theorem 4.5. *It follows for the risk of the estimator* $\widehat{\vartheta}$ *for all* $C \in \mathbf{C}$ *whenever* $\lambda|C| \geqslant 1$*:*

$$\mathbb{E}[(\widehat{\vartheta} - |C|)^2]^{1/2} \leqslant (1 + c\alpha(\lambda, C))\mathbb{E}[(\widehat{\vartheta}_{oracle} - |C|)^2]^{1/2} + r(\lambda, C)\,,$$

with constant $c > 0$ *and* $\alpha(\lambda, C), r(\lambda, C)$ *from Theorem 4.4. For any* $C \in \mathbf{C}$ *and* $\lambda > 0$ *we have* $\alpha(\lambda, C) \leqslant 1 + \frac{1}{\lambda|C|}$*.*

Proof. In view of $\lambda_\circ = \lambda|\widehat{C}|$, we have $\widehat{\vartheta} = \widehat{\vartheta}_{pseudo} - \lambda^{-1}N_\partial \mathrm{e}^{-\lambda|\widehat{C}|}$ and we derive as in (4.4) and (4.6) with some constants $c_1, c_2 > 0$

$$\mathbb{E}[(\widehat{\vartheta} - \widehat{\vartheta}_{pseudo})^2] \leqslant \lambda^{-2}\mathbb{E}[N_\partial^4]^{1/2}\mathbb{E}[\mathrm{e}^{-4\lambda|\widehat{C}|}]^{1/2} \leqslant c_1^2 \exp\big(-2c_2(\lambda|C|)^{(d-1)/(d+1)}\big).$$

To establish the oracle inequality, we apply the triangle inequality in L^2-norm together with Theorems 3.2 and 4.4.

The universal bound on $\alpha(\lambda, C)$ follows from the rough bound $\mathbb{E}[|C \setminus \widehat{C}|^2] \leqslant |C|\mathbb{E}[|C \setminus \widehat{C}|]$. □

Note that the remainder term $r(\lambda, C)$ is exponentially small in $\lambda|C|$. Therefore, an immediate implication of Theorem 4.5 is that asymptotically our estimator $\widehat{\vartheta}$ is minimax rate-optimal in all dimensions, where the lower bound is proved in the next section. Yet, even more is true: the oracle inequality is in all well studied cases *exact* in the sense that $\alpha(\lambda, C) \to 0$ holds for $\lambda \to \infty$ such that the the UMVU risk of $\widehat{\vartheta}_{oracle}$ is attained asymptotically.

Lemma 4.6. *We have tighter bounds on* $\alpha(\lambda, C)$ *from Theorem 4.4 in the following cases:*

1. *for* $d = 1, 2$ *and* $C \in \mathbf{C}$ *arbitrary:* $\alpha(\lambda, C) \lesssim (\lambda|C|)^{-2/(d+1)}$*,*
2. *for* $d \geqslant 2$*,* C *with* C^2*-boundary of positive curvature:* $\alpha(\lambda, C) \lesssim (\lambda|C|)^{-2/(d+1)}$*,*
3. *for* $d \geqslant 2$ *and* C *a polytope:* $\alpha(\lambda, C) \lesssim \lambda^{-1}(\log(\lambda|C|))^{d-1}$*.*

Proof. Let us restrict to $|C| = 1$, the case of general volume follows by rescaling. In view of the expectation upper bound (3.5), the main issue is to bound $\mathrm{Var}(|C \setminus \widehat{C}|)/\mathbb{E}[|C \setminus \widehat{C}|]$ uniformly. Case (1) follows from Pardon (2011), where $\lambda\,\mathrm{Var}(|C \setminus \widehat{C}|) \sim \mathbb{E}[|C \setminus \widehat{C}|]$ is established.

Figure 2: The two convex sets (blue), observations (points), their convex hulls (black lines) and dilated convex hulls (black dashed lines).

For case (2) with smooth boundary, the upper bound for the variance, $\mathrm{Var}(|C\setminus\widehat{C}|) \lesssim \lambda^{-(d+3)/(d+1)}$, was obtained in Reitzner (2005), while the lower bound for the first moment, $\mathbb{E}[|C\setminus\widehat{C}|] \gtrsim \lambda^{-2/(d+1)}$, is due to Schütt (1994).

For the case (3) of polytopes, the upper bound $\mathrm{Var}(|C\setminus\widehat{C}|) \lesssim \lambda^{-2}(\log\lambda)^{d-1}$ was obtained in Bárány and Reitzner (2010), while the lower bound for the first moment, $\mathbb{E}[|C\setminus\widehat{C}|] \gtrsim \lambda^{-1}(\log\lambda)^{d-1}$, was proved in Bárány and Larman (1988). The expectation upper bound from Remark 3.3 thus yields the result. □

One could conjecture that $\lambda\,\mathrm{Var}(|C\setminus\widehat{C}|) \sim \mathbb{E}[|C\setminus\widehat{C}|]$ holds universally for all convex sets in arbitrary dimensions and thus that the oracle inequality is always exact. Proving such a universal bound is a challenging open problem in stochastic geometry, strongly connected to the discussion on universal variance asymptotics in terms of the floating body by Bárány and Reitzner (2010).

5 Finite sample behaviour and dilated hull estimator

In this section, we demonstrate the performance of the main estimator $\widehat{\vartheta}$ numerically and compare it to other estimators including the naive estimator $|\widehat{C}|$, the naive oracle estimator N/λ, the UMVU oracle estimator $\widehat{\vartheta}_{oracle}$ and the plug-in MLE estimator $\widehat{\vartheta}_{plugin} = |\widehat{C}|(1+N_\partial/N)$. The main competitor from the literature is a rate-optimal estimator proposed in Gayraud (1997).



Figure 3: Monte Carlo RMSE estimates for the studied estimators for the volume of two convex sets: a polygon and an ellipse.

In their construction, the whole sample is divided into three equal parts X, X' and X'' of sizes $N^\star$ (without loss of generality $N^\star \in \mathbb{N}$) and the estimator is given by

$$\widehat{\vartheta}_G = |\widehat{C}| + \frac{|\widehat{C}''|}{N^\star} \sum_{i=1}^{N^\star} \mathbf{1}(X_i' \notin \widehat{C}) \,,$$

where $\widehat{C}''$ is the convex hull of the third sample X''. The data points are simulated for two convex sets: an ellipse and a polygon; see Figure 2.

The RMSE estimate normalised by the area of the true set is based on $M = 500$ Monte Carlo iterations in each case. The results of the simulations are depicted in Figure 3 where $n = \lambda|C|$ denotes the expected total number of points. The worst convergence rate of N/λ is clearly visible. More importantly, we see that the RMSE of $\widehat{\vartheta}$ approaches the oracle risk for larger n (i.e. λ) as the oracle inequality predicts. It is also conspicuous that in the studied cases the plug-in estimator $\widehat{\vartheta}_{plugin}$ and the estimator $\widehat{\vartheta}$ perform rather similarly. This is explained by the fact that the number of points N_∂ on the convex hull increases with a moderate speed in the two-dimensional case, $\mathbb{E}[N_\partial] = O(\lambda^{1/3})$, which results in a small difference between the multiplication factors N_∂/N and $N_\partial/(N_\circ + 1)$. The simulations in two dimensions were implemented using the R package "spatstat" by Baddeley and Turner

(2005). To illustrate the sub-optimality of the plug-in estimator $\widehat{\vartheta}_{plugin}$ in high dimensions, we provide results of numerical simulations in dimensions $d = 3, 4, 5, 6$ for the case when the true set C is a unit cube $C = [0, 1]^d$, see Figure 5. The simulations were implemented using the R package "geometry" by Habel, Grasman, Stahel, and Sterratt (2014).

As an application of the obtained results, we propose a new estimator for the convex set itself:

$$\begin{aligned}\widetilde{C} &\stackrel{\text{def}}{=} \left\{\widehat{x}_0 + \Big(\frac{\widehat{\vartheta}}{|\widehat{C}|}\Big)^{1/d}(x - \widehat{x}_0) \,\Big|\, x \in \widehat{C}\right\} \\ &= \left\{\widehat{x}_0 + \Big(\frac{N+1}{N_\circ + 1}\Big)^{1/d}(x - \widehat{x}_0) \,\Big|\, x \in \widehat{C}\right\},\end{aligned}$$

which is just the dilation of the convex hull $\widehat{C}$ from its barycentre $\widehat{x}_0$, see the dashed polygons in Figure 2. Since the convex hull is a sufficient statistic (for known λ), the points in its interior do not bear any information on the shape of C itself such that the barycentre is a reasonable choice. There are, of course, other enlargements of the convex hull conceivable like $\operatorname{argmin}_{B\in\mathbf{C},|B|=\widehat{\vartheta}} d_H(B, \widehat{C})$, the convex set closest (in Hausdorff distance) to $\widehat{C}$ with volume $\widehat{\vartheta}$. The intuition behind these estimators is based on the observation that once the volume of the true set is known, we can estimate the set itself faster (in the constant), and $\widehat{\vartheta}$ is a reasonable substitute for the true volume due to its fast rate of convergence.

A detailed analysis is not pursued here, but in a small simulation study we investigate the behaviour of the new dilated hull estimator for the above polygon. The error ratio $\mathbb{E}[|C\Delta\widehat{C}|]/\mathbb{E}[|C\Delta\widetilde{C}|]$ in terms of the symmetric difference $A\Delta B = (A\backslash B)\cup(B\backslash A)$ is approximated in $M = 500$ Monte Carlo iterations and shown in Figure 4. It turns out that the dilation significantly improves the convex hull as an estimator for C, especially for a small number of observations.

6 Appendix

6.1 Proof of Theorem 3.1

The proof is split into several statements, which might be of interest on their own.



Figure 4: Monte Carlo error ratio for the convex hull and its dilation when the true set is a polygon.

Lemma 6.1. *The random variable* $\mathcal{N}(\mathcal{K})$ *is measurable with respect to* $\mathcal{F}_{\mathcal{K}}$ *for any stopping set* $\mathcal{K}$*.*

Proof. The proof is just a generalisation of the analogous statement for time-indexed stochastic processes, see e.g. Proposition 2.18 in Karatzas and Shreve (2012). For this, the notions are extended to the partial order $\subseteq$ and then the right-continuity of $(\mathcal{N}(K), K \in \mathbf{K})$ (with respect to inclusion) implies its progressive measurability and thus in turn the measurability of $\mathcal{N}(\mathcal{K})$. □

Next, observe that the set-indexed process $(\mathcal{N}(K), K \in \mathbf{K})$ has independent increments, i.e. for $K_1, \dots, K_m \in \mathbf{K}$ with $K_i \subseteq K_{i+1}$, $i = 1, \dots, m-1$, the random variables $\mathcal{N}(K_{i+1}) - \mathcal{N}(K_i) = \mathcal{N}(K_{i+1} \setminus K_i)$ are independent (by the independence of the PPP on disjoint sets). In fact, we show in Proposition 6.2 that the process $\mathcal{N}$ is even a strong Markov process. In addition, Proposition 6.2 yields (3.1) using that the closed complement $\widehat{\mathcal{K}} = \overline{\widehat{C}^c}$ of the convex hull is a stopping set.

Proposition 6.2. *The set-indexed process* $(\mathcal{N}(K), K \in \mathbf{K})$ *is strong Markov at every stopping set* $\mathcal{K}$*. More precisely, conditionally on* $\mathcal{F}_{\mathcal{K}}$ *the process* $(\mathcal{N}(K \setminus \mathcal{K}), K \in \mathbf{K})$ *is a Poisson point process with intensity* λ *on* $\mathcal{K}^c$*. In particular,* $\mathcal{N}(K \setminus \mathcal{K}) \,\big|\, \mathcal{F}_{\mathcal{K}} \sim \mathit{Poiss}(\lambda |K \setminus \mathcal{K}|)$ *holds for all* $K \in \mathbf{K}$*.*

Remark 6.3. *The fact that the increments* $\mathcal{N}(\mathcal{K} \cup K) - \mathcal{N}(\mathcal{K})$ *are independent of* $\mathcal{F}_{\mathcal{K}}$ *can be derived from a general theorem about the strong Markov*

property for random fields in Thm. 4 in Rozanov (1982). See also Zuyev (2006) for a discussion of the strong Markov property and its applications in stochastic geometry. These statements, however, do not provide a distributional characterisation of the increments of the process.

Proof. A set-indexed, $(\mathcal{F}_K)$-adapted integrable process $(X_K, K \in \mathbf{K})$ is called a martingale if $\mathbb{E}[X_B|\mathcal{F}_A] = X_A$ holds for any $A, B \in \mathbf{K}$ with $A \subseteq B$. By the independence of increments, the process $M_K \stackrel{\text{def}}{=} \mathcal{N}(K) - \lambda|K|$, $K \in \mathbf{K}$, is clearly a martingale with respect to its natural filtration $(\mathcal{F}_K, K \in \mathbf{K})$. Then also the process

$$\widetilde{M}_K \stackrel{\text{def}}{=} M_{K\cup\mathcal{K}} - M_{\mathcal{K}} = \mathcal{N}(K \setminus \mathcal{K}) - \lambda|K \setminus \mathcal{K}|$$

is a martingale with respect to the filtration $\widetilde{\mathcal{F}}_K \stackrel{\text{def}}{=} \mathcal{F}_K \vee \mathcal{F}_{\mathcal{K}} = \mathcal{F}_{K\cup\mathcal{K}}$ because for $K_1, K_2 \in \mathbf{K}$ with $K_1 \subseteq K_2$ the optional sampling theorem (see e.g. Zuyev (1999)) yields

$$\mathbb{E}[\widetilde{M}_{K_2}|\widetilde{\mathcal{F}}_{K_1}] = \mathbb{E}[M_{K_2\cup\mathcal{K}} - M_{\mathcal{K}}|\mathcal{F}_{K_1\cup\mathcal{K}}] = M_{K_1\cup\mathcal{K}} - M_{\mathcal{K}} = \widetilde{M}_{K_1},$$

noting that $K_1 \cup \mathcal{K}$ is again a stopping set.

This implies that $\lambda|K\setminus\mathcal{K}|$, conditionally on $\mathcal{K}$, is the deterministic compensator of the process $\widetilde{N}_K = \mathcal{N}(K \setminus \mathcal{K})$. Then, due to the martingale characterisation of the set-indexed Poisson process, see Thm. 3.1 in Ivanoff and Merzbach (1994) (analogue of Watanabe's characterisation for the Poisson process), the process $\widetilde{N}_K$, conditionally on $\mathcal{F}_{\mathcal{K}}$, is a Poisson point process with mean measure $\widetilde{\mu}(A) = \lambda|A \cap \mathcal{K}^c|$. □

The last statement of Theorem 3.1, that $\mathcal{F}_{\widehat{\mathcal{K}}} = \sigma(\widehat{C})$ is shown next. It can be seen as a generalisation of the interesting fact that for a time-indexed Poisson process the sigma-algebra $\sigma(\tau)$ associated with the first jump time τ coincides with the sigma-algebra of τ-history $\mathcal{F}_\tau$.

Lemma 6.4. *The sigma-algebra $\sigma(\widehat{C})$ coincides with the sigma-algebra $\mathcal{F}_{\widehat{\mathcal{K}}}$ of $\widehat{\mathcal{K}}$-history, i.e. $\sigma(\widehat{C}) = \mathcal{F}_{\widehat{\mathcal{K}}}$.*

Proof. Since $\widehat{\mathcal{K}}$ is $\mathcal{F}_{\widehat{\mathcal{K}}}$-measurable by Lemma 1 in Zuyev (1999) and $\widehat{C} = \overline{\widehat{\mathcal{K}}^c}$, it is evident that $\sigma(\widehat{C}) \subseteq \mathcal{F}_{\widehat{\mathcal{K}}}$. The other direction is more involved. We use that the sigma-algebra $\mathcal{F}_{\widehat{\mathcal{K}}}$ coincides with the sigma-algebra $\sigma(\{\mathcal{N}(\widehat{\mathcal{K}} \cap K), K \in$

$\mathbf{K}\}$) generated by the process stopped at $\widehat{\mathcal{K}}$. This statement can be derived from Thm. 6, Ch. 1 in Shiryaev and Aries (2007). Note that their assumption (1.11) is satisfied in our case, because for all $K \in \mathbf{K}$ and $\omega \in \Omega$ there is ω' such that $\mathcal{N}(U \cap K, \omega) = \mathcal{N}(U, \omega')$ for all $U \in \mathbf{K}$, which simply says that observing points in $K \in \mathbf{K}$ there might be no points outside K. Finally, observe that by definition of the convex hull $\mathcal{N}(\overline{\widehat{C}^c} \cap K) = \mathcal{N}((\partial \widehat{C}) \cap K)$. Modulo null sets, $\mathcal{N}((\partial \widehat{C}) \cap K)$ counts the number of vertices of $\widehat{C}$ in K and is thus $\sigma(\widehat{C})$-measurable. □

Proof of Lemma 4.1. Using that the bias of the oracle estimator $\widehat{\vartheta} = |\widehat{C}| + N_\partial/(N_\circ + 1)|\widehat{C}|$ is exponentially small, it remains to compare its expectation with the expectation of the plug-in estimator $\widehat{\vartheta}_{plugin}$ to show (4.1):

$$\begin{aligned}\mathbb{E}[\widehat{\vartheta} - \widehat{\vartheta}_{plugin}] &= \mathbb{E}\Big[|\widehat{C}|\big(\frac{N_\partial}{N_\circ + 1} - \frac{N_\partial}{N}\big)\Big] = \mathbb{E}\Big[|\widehat{C}|\frac{N_\partial^2 - N_\partial}{(N_\circ + 1)(N_\circ + N_\partial)}\Big] \\ &\geqslant \frac{d}{d+1}\mathbb{E}\Big[\frac{|\widehat{C}|N_\partial^2}{(N_\circ + 1)2\lambda|C|}\mathbf{1}(N \leqslant 2\lambda|C|)\Big],\end{aligned}$$

where in the last line we have used $|\widehat{C}| > 0$ only if $N_\partial \geqslant d + 1$ and in this case $N_\partial^2 - N_\partial \geqslant \frac{d}{d+1}N_\partial^2$. Using $\mathbb{E}[(N_\circ + 1)^{-1} \,|\, \widehat{C}] = \lambda_\circ^{-1}(1 - \mathrm{e}^{-\lambda_\circ})$ from above, we obtain after writing $\mathbf{1}(N \leqslant 2\lambda|C|) = 1 - \mathbf{1}(N > 2\lambda|C|)$

$$\begin{aligned}\mathbb{E}[\widehat{\vartheta} - \widehat{\vartheta}_{plugin}] &\geqslant \frac{d}{d+1}\Big(\mathbb{E}\Big[\frac{N_\partial^2|\widehat{C}|(1 - \mathrm{e}^{-\lambda_\circ})}{2\lambda_\circ\lambda|C|}\Big] - \mathbb{E}\Big[\frac{N_\partial^2|\widehat{C}|}{2\lambda|C|}\mathbf{1}(N > 2\lambda|C|)\Big]\Big) \\ &\geqslant \frac{d}{d+1}\Big(\frac{\mathbb{E}[N_\partial^2(1 - \mathrm{e}^{-\lambda_\circ})]}{2\lambda^2|C|} - \frac{\mathbb{E}\big[N^2\mathbf{1}(N > 2\lambda|C|)\big]}{2\lambda}\Big).\end{aligned}$$

By Cauchy-Schwarz inequality and large deviations similarly to (4.4), the first term is bounded from below by a constant multiple of $\mathbb{E}[|C \setminus \widehat{C}|]^2/|C|$ in view of $\mathbb{E}[N_\partial^2] \geqslant \lambda^2\mathbb{E}[|C \setminus \widehat{C}|]^2$, see e.g. Section 2.3.2 in Brunel (2014). Because of $N \sim \mathrm{Poiss}(\lambda|C|)$, the second term is of order $\lambda|C|^2\mathrm{e}^{-\lambda|C|}$ and thus asymptotically of much smaller order. □

References

Baddeley, A., and R. Turner (2005): "spatstat: An R Package for Analyzing Spatial Point Patterns," *Journal of Statistical Software*, 12(6), 1–42, http://www.jstatsoft.org/v12/i06/.

Bárány, I., and D. Larman (1988): "Convex bodies, economic cap coverings, random polytopes," *Mathematika*, 35(2), 274–291.

Bárány, I., and M. Reitzner (2010): "On the variance of random polytopes," *Advances in Mathematics*, 225(4), 1986–2001.

Beermann, M., and M. Reitzner (2015): "Beyond the Efron-Buchta identities: Distributional Results for Poisson Polytopes," *Discrete & Computational Geometry*, 53(1), 226–244.

Brunel, V.-E. (2013): "A universal deviation inequality for random polytopes," arXiv:1311.2902.

——— (2014): "Non-parametric estimation of convex bodies and convex polytopes," Theses, Université Pierre et Marie Curie - Paris VI ; University of Haifa, https://tel.archives-ouvertes.fr/tel-01066977.

Davis, R., E. Mulrow, and S. Resnick (1987): "The convex hull of a random sample in $\mathbb{R}^2$," *Stochastic Models*, 3(1), 1–27.

Dwyer, R. (1988): "On the convex hull of random points in a polytope," *Journal of Applied Probability*, 25(4), 688–699.

Gayraud, G. (1997): "Estimation of functionals of density support," *Mathematical Methods of Statistics*, 6(1), 26–46.

Guntuboyina, A. (2012): "Optimal rates of convergence for convex set estimation from support functions," *The Annals of Statistics*, 40(1), 385–411.

Habel, K., R. Grasman, A. Stahel, and D. Sterratt (2014): *geometry: Mesh generation and surface tesselation.*R package version 0.3-5, http://CRAN.R-project.org/package=geometry.

Ivanoff, B., and E. Merzbach (1994): "A martingale characterization of the set-indexed Poisson process," *Stochastics: An International Journal of Probability and Stochastic Processes*, 51(1-2), 69–82.

Karatzas, I., and S. Shreve (2012): *Brownian motion and stochastic calculus*. Springer.

Karr, A. (1991): *Point processes and their statistical inference*, Probability: Pure and Applied. CRC press.

Kingman, J. (1992): *Poisson processes*, vol. 3. Oxford University Press.

Korostelev, A., and A. Tsybakov (1993): *Minimax theory of image reconstruction*. Springer.

——— (1994): "Asymptotic efficiency in estimation of a convex set," *Problemy Peredachi Informatsii*, 30(4), 33–44.

Kutoyants, Y. (1998): *Statistical Inference for Spatial Poisson Processes*, Lecture Notes in Statistics. Springer.

Mammen, E., and A. Tsybakov (1995): "Asymptotical Minimax Recovery of Sets with Smooth Boundaries," *The Annals of Statistics*, 23(2), 502–524.

Meister, A., and M. Reiss (2013): "Asymptotic equivalence for nonparametric regression with non-regular errors," *Probability Theory and Related Fields*, 155(1-2), 201–229.

Molchanov, I. (2006): *Theory of random sets*. Springer.

Pardon, J. (2011): "Central limit theorems for random polygons in an arbitrary convex set," *The Annals of Probability*, 39(3), 881–903.

Privault, N. (2012): "Invariance of Poisson measures under random transformations," *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, 48(4), 947–972.

Reiss, M., and L. Selk (2015): "Estimating nonparametric functionals efficiently under one-sided errors," *Bernoulli (to appear)*, arXiv:1407.4229.

Reitzner, M. (2003): "Random polytopes and the Efron–Stein jackknife inequality," *The Annals of Probability*, 31(4), 2136–2166.

——— (2005): "Central limit theorems for random polytopes," *Probability Theory and Related Fields*, 133(4), 483–507.

Resnick, S. (2013): *Extreme values, regular variation and point processes*. Springer.

Rozanov, Y. (1982): *Markov random fields*. Springer.

Schütt, C. (1994): "Random polytopes and affine surface area," *Mathematische Nachrichten*, 170(1), 227–249.

Shiryaev, A., and A. Aries (2007): *Optimal stopping rules*. Springer.

Zuyev, S. (1999): "Stopping sets: Gamma-type results and hitting properties," *Advances in Applied Probability*, 31(2), 355–366.

——— (2006): "Strong Markov property of Poisson processes and Slivnyak formula," in *Case Studies in Spatial Point Process Modeling*, pp. 77–84. Springer.



Figure 5: Monte Carlo RMSE estimates for the studied estimators for the volume of the unit cube $C = [0,1]^d$ in dimensions $d = 3, 4, 5, 6$.